

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- ① BLACK BORDERS
 - TEXT CUT OFF AT TOP, BOTTOM OR SIDES
 - FADED TEXT
 - ILLEGIBLE TEXT
 - SKEWED/SLANTED IMAGES
 - COLORED PHOTOS
- ② BLACK OR VERY BLACK AND WHITE DARK PHOTOS
 - GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵: C12N 15/82, 15/54, 5/10, A01H 5/00, A23L 1/00, 1/09, 1/307, A23K 1/00, C08B 37/00	A1	(11) International Publication Number: WO 94/14970 (43) International Publication Date: 7 July 1994 (07.07.94)
(21) International Application Number: PCT/NL93/00279 (22) International Filing Date: 28 December 1993 (28.12.93) (30) Priority Data: 92204098.5 28 December 1992 (28.12.92) EP (34) Countries for which the regional or international application was filed: NL et al. 9300646 15 April 1993 (15.04.93) NL (71) Applicant (for all designated States except US): STICHT- ING SCHEIKUNDIG ONDERZOEK IN NEDERLAND [NL/NL]; Laan van Nieuw Oost Indië 131, NL-2395 BM The Hague (NL). (72) Inventors; and (75) Inventors/Applicants (for US only): SMEEKENS, Josephus, Christianus, Maria [NL/NL]; Westrevelaan 7, NL-3971 AE Driebergen (NL). EBSKAMP, Michaël, Johannes, Marcus [NL/NL]; Rijksweg 55, NL-3454 JD De Meern (NL). WEISBEEK, Petrus, Jacobus [NL/NL]; Baarnseweg 49, NL- 3734 CA Den Dolder (NL). (74) Agent: HOUTINK, Reinoud; Arnold & Siedsma, Sweelinck- plein 1, NL-2517 GK The Hague (NL).		(81) Designated States: AU, BB, BG, BR, BY, CA, CZ, FI, HU, JP, KP, KR, KZ, LK, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SE, SK, UA, US, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>
(54) Title: METHOD FOR OBTAINING TRANSGENIC PLANTS SHOWING A MODIFIED FRUCTAN PATTERN (57) Abstract <p>The invention relates to a method for obtaining transgenic plants showing a modified fructan pattern as compared to non-transformed plants, comprising the steps of: a) preparing a DNA construct comprising one or more fructosyltransferase genes, or modified versions thereof, being operably linked to a promoter sequence active in plants and a terminator sequence active in plants; b) transforming a plant cell with the construct; and c) regenerating a transgenic plant from the transformed plant cell. Preferably, the 5' untranslated region of the fructosyltransferase gene is modified such that the expression of the fructosyltransferase is not negatively affected. The DNA construct preferably also comprises a targeting sequence upstream from the fructosyltransferase gene for directing the fructosyltransferase to a specific plant tissue or plant cell compartment. The fructosyltransferases produced in the transgenic plants change the fructan pattern of the plants thus leading to different plant performance due to altered sink-source relations and yield, or increased tolerance for drought, cold or other stresses, higher dry matter content, better taste or storability, improved nutritional value. The plants may also suitably be used as raw material for fructan production.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

METHOD FOR OBTAINING TRANSGENIC PLANTS
SHOWING A MODIFIED FRUCTAN PATTERN

The invention relates to a method for obtaining transgenic plants showing a modified fructan pattern as compared to non-transformed plants, a DNA construct for producing transgenic plants or plant tissues, the transgenic plants or plant tissues showing a modified fructan pattern as compared to non-transformed plants, the fructans isolated from the plants or tissues, the use of the transgenic plants, tissues and fructans for various applications and transgenic seed.

Plants produce many carbohydrates useful for food and non-food applications. Important examples of such carbohydrates are starch, cellulose and sucrose. These compounds are used by man either as such or in a modified form for food and industrial purposes. Crop species which produce these carbohydrates have been obtained by traditional plant breeding methods. Next to the above mentioned well known carbohydrates many other plant carbohydrates with properties useful to man can be present in plants.

Plant carbohydrates can be divided into two groups depending on their function for the plants. The first group comprises structural carbohydrates which are usually part of the extracellular matrix. The most prominent member of this group is cellulose. The second group are the non-structural storage carbohydrate which can serve as a long term or short term (transitory) carbohydrate stores. Examples of such carbohydrates are starch, sucrose and fructans.

Fructan is a polymer consisting mostly of repeating fructose units. Fructans are mostly found in plants species which are not adapted to economic farming practices. Fructans occur in monocotyledons (esp. Poaceae, Liliaceae) and in dicotyledons (esp. Compositae) (Pollock and Cairns 1991, Ann. Rev. Plant Physiol. Plant Mol. Biol. 42, 77-101; Hendry 1987, New Phytol. 106, 201-216).

Next to its role as a plant carbohydrate reserve other functions have been proposed for fructans. These

functions include for example tolerance to dry and cold climates (cold induced desiccation) (Pontis 1989, J. Plant Physiol. 134, 148-150; Gonzales et al. 1990 New Phytol. 115, 319-323).

5 However, the fructans produced by plants usually have limited functionality. For these reasons fructans have as of yet not found great application in food and non-food products in spite of the recognized great commercial potential (Fuchs 1991 Biochem. Soc. Transact. 19, 555-560, A. Fuchs, ed. Inulin and Inulin-containing Crops, Elsevier, 1993) especially when compared to other carbohydrates.

10 One of the major problems which limit the usefulness of plant fructans is the limited range of plant species in which fructans accumulate to a substantial level. In many important current crop plants, among which sugar beet and potato, fructans are either absent or found at very low levels. Plants which do store fructans to some extent often have unfavourable agronomic properties. An example of such a plant is Helianthus tuberosus.

20 Another problem is the limited functionality of the fructans produced. The functionality is inter alia determined by the length of the fructan chains which in plants seldomly exceeds a so-called "Degree of Polymerization" (DP) of the monosaccharides of 100. Usually much smaller fructans are found and often the DP of these fructans decreases dramatically prior to harvest or following harvest and/or storage. Low DP's severely limit the usefulness of fructans in subsequent processes and can result in reduced yields.

30 It has now been found that it is possible to transform certain plants in order to provide for expression of fructans in suitable host plants.

 The invention thus provides a method for obtaining transgenic plants showing a modified fructan pattern as compared to the non-transformed plants, comprising the steps of:

35 a) preparing a DNA construct comprising one or more fructosyltransferase genes, or modified versions thereof, which in plants seldomly exceeds a so-called "Degree of Polymerization" (DP) of the monosaccharides of 100. Usually much smaller

of, being operably linked to a promotor sequence active in plants and a terminator sequence active in plants;

b) transforming a plant cell with the construct;

and

5 c) regenerating a transgenic plant from the transformed plant cell.

Fructosyltransferase genes are genes encoding enzymes which catalyze the production of fructose-fructose linkage. It has been proposed in the international patent application WO89/12386 to produce transgenic plants comprising a foreign gene encoding an enzyme capable of polymerizing a carbohydrate, such as for example various sucrases, in order to modify the soluble solids composition of the plant cells. WO89/12386 discloses inter alia the use of 10 dextran sucrose. This enzyme uses sucrose as a source for the polymerization of glucose units. The present invention however relates to the polymerization of fructose units.

The present inventors found that by the presence of the 5' microbial untranslated region in the fructosyltransferase gene SacB, in which one out of frame ATG codon precedes the initiator ATG codon it is not possible to obtain acceptable expression levels of the said gene. It was thus found according to another aspect of the invention that suitable modification of the 5' untranslated region of a 15 fructosyltransferase gene increases the expression level of the fructosyltransferase gene in the plant cell. A suitable modification comprises any modification by which the expression level of the fructosyltransferase gene is not negatively affected. In practice this means that any sequence negatively affecting the expression level is to be removed. 20 Preferably the 5' untranslated region up to the ATG codon is entirely removed from the microbial gene.

With the term "fructan pattern" is meant the distribution of the fructan in the different plant tissues and plant cell compartments such as cytosol, vacuole, apoplast etc.. Some plants naturally do produce fructans whereas others do not. The fructan pattern of a plant species not producing fructans in the non-transformed plant may be 25

changed by the introduction of a gene encoding a fructosyltransferase. The fructan distribution in a plant may also be changed by the redirection of the metabolic flow in plants or plant cells to certain plant or plant cell compartments.

5 Sucrose which is the substrate for the fructosyltransferases is a carbohydrate which is found in several different locations. It is synthesized in the cytoplasm and significant quantities can be found in cytosol, vacuole and the extracellular space (the apoplast).


10 Since biochemical processes in plants cells are also often limited to a single or a few cellular compartments it is desirable to promote the accumulation of the products of the newly introduced genes into a specific compartment. Therefore targeting sequences which are specific for cellular compartments must be present on the fructosyltransferases which are expressed in the transgenic plants. Specific amino acid regions for targeting to the different cellular locations have been identified and analyzed. The DNA sequences encoding these targeting regions can be coupled to the genes of interest in such a way that upon expression of the chimeric gene a protein is produced in which the targeting information is recognized and used by the cell for correct targeting to the cellular location of interest (Oakes et al. EPO 0486683, WO 91/19808, Symons et al. Bio/Technology 8, 217-221 (1990), Pen J., et al. Bio/Technology 10, 292-296 (1992)).

 In a preferred embodiment of the invention the expression cassette therefore also comprises a targeting sequence for directing the fructosyltransferases to one or more specific plant or plant cell compartments. The targeting regions may be any DNA sequence having the ability of directing the fructosyltransferases to one or more of these specific plant or plant cell compartments. Examples of preferred targeting regions are the signal sequence and vacuolar targeting sequence of the carboxypeptidase Y (cpv) gene, or the signal sequence and apoplasmatic targeting sequence of the pathogenesis-related protein S gene (pr-s). By adding the targeting sequence to the DNA construct the

present invention provides a method for producing transgenic plants in which the fructosyltransferase may be directed to one or more specific cell compartments thus leading to a desired fructan pattern in the plant or plant cell.

Often it is beneficial to the plant to control not only the location but also the timing of expression of the introduced genes. For example it is usually desired to limit the expression of the newly introduced enzymatic activities to specific parts of the plants, e.g. harvestable organs like tubers, fruits or seeds. Moreover it is often desired to initiate expression in these organs at a certain developmental stage. This is certainly the case when the expression of the introduced genes interferes with normal development of such organs.

The fructosyltransferases of the present invention use sucrose as a substrate to synthesize high molecular weight fructans. Many micro-organisms contain fructosyltransferases which have the ability to produce fructans (often called levans) from sucrose (A. Fuchs 1959, Thesis, University of Leiden, Han 1989 Adv. Appl. Microbiol. 35, 171-194). These enzymes can transfer fructose moieties from sucrose to a fructan acceptor molecule to produce high molecular weight fructans. Since these fructan acceptor molecules are originally derived from sucrose they may still contain a terminal glucose molecule. The reaction mechanism is as follows (review see Han 1989 Adv. Appl. Microbiol. 35, 171-194):



G-F = sucrose, G = glucose, F = fructose

The glycosidic linkage which interconnects the fructose units can be a (2-1)- or a (2-6)- type depending on the particular fructosyltransferase. In micro-organisms both linkage types can be found in a single fructan molecule. Functionality of a fructan molecule depends on the type of

backbone ((2-1)- or (2-6)) and the degree of branching and the degree of polymerization.

Many micro-organisms, such as bacteria, yeasts, fungi etc. contain fructosyltransferases which synthesize high DP fructans from sucrose. Usually the fructans are deposited outside the cell.

Two examples of fructosyltransferase producing bacteria are Bacillus subtilis (Steinmetz et al. 1985 Mol. Gen. Genet. 200, 220-228) and Streptococcus mutans (Shiroza, and Kuramitsu 1988 J. Bacteriol. 170, 810-816. In B. subtilis the sacB gene encodes the structural gene for levansucrase, a fructosyltransferase. This enzyme can convert sucrose into a fructose polymer with DP's which can easily exceed 10000. The linkage type found in this fructan produced by the levansucrase is mainly of the (2-6)-type with extensive (2-1)-branching.

In another micro-organism S. mutans the ftf gene encodes a fructosyltransferase which can also produce fructans of very high DP. The fructose units in this fructan are mainly (2-1)-linked type with (2-6)-branching (Rosell and Birkhed 1974 Acta Chem. Scand. B28, 589).

Bacterial fructosyltransferases have a relatively low K_m for sucrose, approx. 20 mM. Sucrose concentrations in most plant is considerably higher and therefore these enzymes should be active in plants. Another important property of bacterial fructosyltransferases is their ability to synthesize levans at low temperatures up to 0°C (Tanaka et al. 1978 Agric. Biol. Chem. 42, 323-326. Plants often encounter such temperatures but in such conditions these enzymes could still be active.

The preferred structural fructosyltransferase gene(s) largely depend on the requirements for degree of polymerisation of the polymer and its degree of branching of the fructans for the various uses.

The present invention is not limited to microbial fructosyltransferases but it is also possible to use fructosyltransferases originating from plants or any other prokaryotic or eukaryotic source.

In plants fructan biosynthesis and degradation has been investigated especially in Helianthus tuberosus and in several grasses (Pollock and Cairns 1991, Ann. Rev. Plant Physiol. Plant Mol. Biol. 42, 77-101; Pollock 1986, New Phytol. 104, 1-24). In different plant families different fructans types are synthesized, linear or branched, with 2-1 and 2-6 glycosidic linkages between the fructose molecules. In plants fructans are stored in the vacuoles. The substrate for fructan biosynthesis is sucrose, however the affinity of these enzymes for sucrose is lower (approx. Km 100) than is the case with the bacterial enzymes. The plant enzymes involved in fructan metabolism are very cold tolerant (Pontis 1989, J. Plant Physiol. 134, 148-150; Gonzales et al. 1990 New Phytol. 115, 319-323) and may therefore be very useful in certain applications.

Nearly all plants are capable of sucrose synthesis which takes place in the cytosol. Sucrose is the major carbohydrate translocation sugar and it is transported from source (net carbohydrate exporting tissue) to sink (net carbohydrate importing tissue) via the vascular system. This transport often includes transfer of sucrose through the extracellular space, the apoplast. Moreover sucrose can also be used by plants for carbohydrate storage. In the latter case it usually accumulates in the vacuoles of the plant cells, e.g. in the roots of sugar beet plants. In conclusion there are two important cellular locations where sucrose is found in plant cells: the cytosol and the vacuole. In addition substantial amounts of sucrose is present in the extracellular space (the apoplast) of many plants.

The preferred targeting region should target the fructosyltransferase to compartments where substrate is available being vacuoles, cytosol and the apoplast, respectively. For high fructan accumulation for instance the vacuole is the preferred compartment. For other purposes (e.g. drought and cold resistance) fructan accumulation in the cytosol or apoplast may be preferable.

Especially the accumulation of sucrose in the vacuoles of many plants makes this cellular location ideally

suited for the induction of fructan biosynthesis by microbial or plant enzymes.

A preferred embodiment of the invention therefore provides a method for the vacuolar expression of microbial, plant or other fructosyltransferases in transgenic plants. This method involves the fusion of DNA sequences which encode plant specific expression signals, a vacuolar localization signal and microbial, plant or other fructosyltransferases.

Two other compartments in which sucrose is available, the cytosol and the apoplast, can also be important sites for fructan accumulation. This can be achieved by the direction of microbial, plant or other fructosyltransferases to the cytosol or apoplast. For cytosolic expression no cellular routing information is required; expression of the mature enzymes is sufficient. For apoplastic expression of fructosyltransferases a targeting region must be added to the enzyme which targets the enzyme to the apoplast.

For expression in plants of the modified fructosyltransferase gene promoter signals must be added to the DNA construct. Such expression promoters can be specific for a certain cell type or can be active in a wide array of cell types. In addition, the timing of expression can be determined by the use of for example developmental-specific promoters. One commonly used promoter for gene expression in plants is the 35S Cauliflower Mosaic Virus promoter which is active in many cell types in the plant independent of the developmental state of the plant.

The preferred promoter may be a tissue-specific or constitutive promoter, strong or weak, depending on target plant and objective. For fructan production in storage organs the preferred promoter would be a sink-specific, strong promoter.

To further improve transcription levels the promoter is often modified to contain an enhancer duplication.

The translation of the mRNA's may be improved by adding a translational enhancer such as the Alfalfa Mosaic

Virus RNA4 translational enhancer signal which must be present in the transcribed 5' untranslated region.

For proper termination of transcription a plant specific terminator sequence must be added to the constructs. An example of such a sequence is the nopaline synthase gene termination sequence.

The present invention is not limited to naturally occurring fructosyltransferases but may equally well be performed by using modified versions thereof. Modifications may influence the activity of the fructosyltransferases in such a way that e.g. the degree of polymerization or the structure of the fructan produced is altered.

The induced accumulation of fructans in transgenic plants using the principle described above will allow for the extraction of fructans from these plants for the purpose of fructan production. Fructans can be accumulated in these plants e.g. in harvestable organs such as roots, beets, leaves, stems, tubers, fruits and seeds.

Genetically modified crop plants which incorporate the fructosyltransferase-encoding constructs mentioned above will allow for the efficient production of a high quality carbohydrate polymer useful to man.

The invention provides a method which provides plants with new biosynthetic capacities which enable them to produce and accumulate fructan. Such plants have improved performance due to altered sink-source relations and yield, or improved performance under abiotic and biotic stresses. Such stresses include but are not limited to drought, light, temperatures, diseases and pests. Improved performance under normal or stress conditions include, but are not limited to, higher dry matter content, better taste or storability, improved nutritional value, etc. Such plants can be suitably used as raw material for fructan production.

According to the present invention a single fructosyltransferase gene or a combination of fructosyltransferase genes of either prokaryotic or eukaryotic origin may be used. These genes can encode enzymes for the biosynthesis of a wide range of fructans.

The present invention further relates to seeds, cuttings, tubers, bulbs, or other parts of the transgenic plants which are useful for the continuous production of further generations of said plants.

5 The fructans produced using transgenic plants of the present invention may be used in various food and non-food applications. Examples include but are not limited to human and animal food products, in the production of fructose syrups, in the production of chemicals and plastics
10 either as such or in a modified form.

The accompanying figures illustrate the present invention wherein

fig. 1 is a summary of the constructs;
fig. 2 shows the construction of plasmids pPA2 and
15 pPB1;

fig. 3 shows the construction of 35S-cpy-sacB-NOS in binary vector (pKP);

fig. 4 shows the construction of 35S-cpy-ftf-NOS in binary vector (pTP);

20 fig. 5 shows the construction of 35S-pr-s-sacB-NOS in binary vector (pKT);

fig. 6 shows the construction of 35S-pr-s-ftf-NOS in binary vector (pTT);

25 fig. 7 shows the construction of 35S-sacB-NOS in binary vector (pKZ);

fig. 8 shows the construction of 35S-ftf-NOS in binary vector (pTZ); and

fig. 9 is an example of a partial acid hydrolysis of fructan isolated from a transgenic KP tobacco plant. This
30 partial hydrolysis pattern is identical to that of fructans produced by Bacillus subtilis. F=fructose, S=sucrose (disaccharide), I=1-kestose (trisaccharide) DP4, 5 and 6=tetra-penta and hexasaccharides, respectively, H=Helianthus tuberosus tuber extract used as standard.

35

The present invention will be illustrated in the following examples which are in no way intended to limit the scope of the present invention.

EXAMPLE 1.**Expression of sacB gene in the vacuole.****1. Selection of genes to be used.**

To create fructan-producing transgenic plants,
5 genes encoding proteins capable of producing fructans were selected. One of these genes, levansucrase, encoded by the sacB gene of Bacillus subtilis (Steinmetz M. et al. Mol. Gen. Genet. 200:220-228(1985)), was used. This enzyme produces mainly branched fructans of the (2-6)-linkage type in
10 the presence of sucrose.

Since sucrose in plant cells can accumulate in vacuoles, this is a preferred site for fructan production. To direct the levansucrase to the vacuole the targeting region of carboxypeptidase Y (Valls L.A. et al. Cell 48:
15 887-897 (1987)) was selected.

2. Construction of 35S-cpy-sacB-NOS in binary vector (pKP).

Plasmid pLS302 has been described by Steinmetz M. et al., supra. The levansucrase (sacB) gene was cloned from
20 this plasmid as a BamHI and PstI fragment into the multiple cloning site of pEMBL9 (Dente L. et al. Nucleic Acids Res. 11, 1645-1655 (1983)) in the corresponding BamHI and PstI sites resulting in plasmid pKA16. General DNA cloning methodology has been described (Sambrook J. et al. Cold Spring
25 Harbour, NY, Cold Spring Harbour Laboratory (1989)).

To create a NcoI site near the mature processing site of the levansucrase gene at nucleotide position 550 (Steinmetz M. et al. supra) site directed mutagenesis as described by Kramer W. et al. (Nucleic Acids Res. 12,
30 9441-9456 (1984)) was performed with the following oligonucleotide 5'-GCAACTCAAGCCATGGCGAAAGAAACG3-' resulting in plasmid pKD22.

At the amino acid position -2 relative to the mature processing site (nucleotide position 544) a phenylalanine was hereby changed to a methionine. The NcoI-PstI
35 fragment, in which the sequence encoding the mature levansucrase protein is present, was used for further cloning.

Plasmid pTS3 containing the carboxypeptidase Y (cpy) gene has been described by Valls L.A. et al., supra.

The first part of the cpy gene was cloned from this plasmid as a ClaI (-695) and BamHI (462) fragment into the multiple cloning site of pEMBL8 (Dente L. et al., supra), in the corresponding AccI and BamHI site resulting in plasmid pCA1.

To create a NcoI site near the mature processing site downstream of the vacuolar targeting sequences of the cpy gene at nucleotide position 330 (Valls, L.A. et al. supra), site directed mutagenesis as described by Kramer W. et al. (supra) was performed with the following oligonucleotide: 5'-CGTGTCAACAAGACCATGGACCCTAA-3'. The resulting plasmid pCB50 contains the first part of the cpy gene with a NcoI site at the mature processing site. At the amino acid position +2 relative to the mature processing site at nucleotide position 334 a isoleucine was replaced by a threonine. At amino acid position +3 relative to the mature processing site at nucleotide position 337 a lysine was replaced by a methionine. The HindIII-NcoI fragment in which the sequence encoding the vacuolar targeting part of carboxypeptidase Y protein is present, was used for further cloning.

The vacuolar targeting sequence of the cpy gene was fused to the mature sequence of the levansucrase gene using a threepoint ligation (Sambrook J. et al. supra).

The following fragments were used in equimolar concentrations:

The HindIII-NcoI fragment of the cpy gene (encoding the vacuolar targeting sequence), the NcoI-PstI fragment of the sacB gene and the fragment containing the vector of pEMBL8 which was digested with HindIII and PstI. The resulting plasmid pKK3 encodes the in frame construct of the carboxypeptidase Y-levansucrase fusion. The correct reading frame of the fusion gene was confirmed by sequence analysis.

The plasmid pMOG18 which contains a plant-specific 35S promoter with enhancer duplication and sequences which enhance translation of the mRNA has been described by Symons et al., Bio/Technology 8, 217-221 (1990)). It contains the

35S-promotor/uidA-gene/NOS-terminator construct. A pBlue-script II SK (Stratagene, San Diego CA) from which the internal BamHI site had been removed, by cutting with BamHI and filling in of the sticky ends with Klenow and religation was used for further cloning. The 35S-uidA-NOS fragment was obtained by digestion with EcoRI and HindIII of pMOG18 and was cloned in this BamHI-pBluescript into the corresponding EcoRI and HindIII site resulting in plasmid pPA2. This plasmid was digested with NcoI and BamHI to remove the uidA gene. The resulting vector was treated with S1-nuclease to remove overhanging ends and then dephosphorylated. From plasmid pKK3 the fragment containing the cpy-sacB fusion was isolated by digestion with AccI and PstI. These sites were blunted with Klenow. These two fragments were ligated resulting in plasmid pKM5. pKM5 contains the 35S-promotor/cpy-sacB-gene-fusion/NOS-terminator construct.

This plasmid, pKM5, was digested first with XhoI and then partially digested with XbaI. The fragment containing the complete construct (35S-cpy-sacB-NOS) was cloned into the XbaI and XhoI site of pMOG23 (Symons et al. supra), a binary plant vector pBIN19 derivative (Bevan M., Nucl. Acids Res. 12: 8711-8721) resulting in plasmid pKP.

3. Transformation of pKP to tobacco plants.

The pKP plasmid was conjugated into Agrobacterium tumefaciens LB4404 (Hoekema A. et al., Nature 303, 179-180 (1983)) in a triparental mating event using the helper plasmid pRK2013 (Lam S.T. et al., Plasmid 13, 200-204 (1985)). The construct was introduced into Nicotiana tabacum var. Petit Havanna (SR1) using the leaf disc transformation method (Horsch R.B. et al., Science 227, 1229-1232 (1985)). Regenerated plants, named KP plants, were selected for kanamycin resistance and grown on MS medium (Murashige and Skoog Physiol. Plant. 15, 473-497 (1962)) containing glucose instead of sucrose. Thereafter plants were grown on soil in the greenhouse and analysed.

4. Analysis of KP plants.

Plants were grown in the greenhouse and leaf material was cut off and ground in an eppendorf tube. Following centrifugation (2' 16000 rpm) 1 μ l of supernatant was spotted on TLC (Cairns A.J. and Pollock C.J. New Phytol. 109, 399-405 (1988)). The TLC was developed three times in 85:15 acetone:water and then treated with urea-spray as described (Wise C.S. et al. Analytical Chemistry 27, 33-36 (1955)). This method preferentially stains fructose and fructose-containing polymers.

Fructose polymers were never detected in non-transformed plants nor in plants transformed with unrelated constructs.

Screening of the transformants, using this method, showed extensive accumulation of fructans in these plants. Fructan expression levels varied between individual plants transformed with the same construct as is normally found in plant transformation experiments. This variation of expression levels depends mainly on the genomic integration site (position effect).

The fructans accumulating in these plants were further studied by isolation of larger quantities (Livingstone III D.A., Plant Physiol. 92, 767-769 (1990)). This fructan was analysed by sizing on a FPLC Superose 6HR 10/30 column (Pharmacia) and fructans were detected in the void volume indicating a degree of polymerisation of >25,000 fructose units. Fructan produced by Bacillus subtilis levansucrase similarly elutes in the void volume of the Superose column.

Partial fructan degradation by acid degradation (See Fig. 9) shows a characteristic pattern of hydrolysis products. The purified plant and bacterial fructans show identical degradation patterns on TLC. Full acid hydrolysis followed by HPLC analysis on an Aminex HPX87C column (Bio-rad) at 85°C using water as eluent shows this fructan to be composed of fructose.

The purified fructans were also analysed by proton-NMR. Fructan isolated from transgenic plants showed no

differences in peak pattern when compared with fructan synthesised by levansucrase from Bacillus subtilis.

We have not found any fructan hydrolysing activity in several plant species (tobacco, sugar beet, tomato, potato). When plant protein extracts are incubated with fructan for prolonged periods of time in phosphate-citrate buffer at pH 5.0 no fructose is released.

Upon senescence plant fructans remain present in the transgenic plants, confirming the stability seen with protein extracts. Fructans accumulate throughout the plant which is in accordance with the unspecific 35S promoter activity in transgenic plants.

Transgenic plants set seed normally. Seed germination is identical to untransformed plants and the construct is stably inherited by following generation which similarly produce fructans.

EXAMPLE 2.

Expression of ftf gene in the vacuole

1. Selection of the genes.

To create fructan-producing plants, genes encoding proteins capable of producing fructans were selected. Another gene, fructosyltransferase encoded by the ftf gene of Streptococcus mutans (Shiroza T. and Kuramitsu H.K., J. Bacteriol. 170, 810-816 (1988)), was used. This enzyme produces mainly branched fructans of the (2-1)-linkage type in the presence of sucrose. Since sucrose in plant cells can accumulate in vacuoles, this is a preferred site for fructan accumulation. A gene encoding a protein capable of targeting itself to the vacuole was selected: carboxypeptidase Y (Valls L.A. et al., supra). From this gene the signal sequence and vacuolar targeting sequence were used as described in Example 1.

2. Construction of 35S-cpv-ftf-NOS in binary vector (pTP)

Plasmid pTS102 has been described by Shiroza T. and Kuramitsu H.K. (supra). The fructosyltransferase (ftf) gene was cloned from this plasmid as an EcoRV and BglII

fragment into the multiple cloning site of pEMBL9 (Dente L. et al., supra) in the compatible SmaI and BamHI sites resulting in plasmid pTA12.

To create a NcoI site near the mature processing site of the ftf gene (nucleotide position 783 (Shiroza T. and Kuramitsu H.K. (supra)) site directed mutagenesis as described by Kramer W. et al. (supra) was performed with the following oligonucleotide:

5'-GGCTCTCTTCTGTTCCATGGCAGATGAAGC-3' resulting in plasmid pTD2.

At amino acid position +1 (nucleotide position 783) relative to the mature processing site a glutamine was hereby changed to a methionine. The NcoI-PstI fragment, in which the sequence encoding the mature fructosyltransferase protein is present, was used for further cloning.

From plasmid pCB50 (as described in Example 1) the HindIII-NcoI fragment in which the sequence encoding the vacuolar targeting part of carboxypeptidase Y protein is present, was used for further cloning.

The vacuolar targeting sequence of the cpy gene was fused to the mature sequence of the fructosyltransferase gene using a threepoint ligation (Sambrook J. et al., supra). The following fragments were used in equimolar concentrations:

The HindIII-NcoI fragment of the cpy gene (encoding the vacuolar targeting sequence), the NcoI-PstI fragment of the ftf gene and the fragment containing the vector of pEMBL8 which was digested with HindIII and PstI. The resulting plasmid pTK1 encodes the in frame construct of the carboxypeptidase Y fructosyltransferase fusion. The fusion is in the correct reading frame which was confirmed by sequence analyses.

Plasmid pPA2 describe in Example 1 was digested with NcoI and BamHI to remove the uidA gene. The resulting vector was treated with S1-nuclease to remove overhanging ends and then dephosphorylated. From plasmid pTK1 the fragment containing the cpy-ftf fusion was isolated by digestion

with AccI and PstI. These sites were blunted with Klenow. These two fragments were ligated resulting in plasmid pTM4. pTM4 contains the 35S-promotor/cpy-ftf-gene-fusion/NOS-terminator construct.

5 This pTM4 was digested with XhoI and then partially digested with XbaI. The fragment containing the complete construct (35S-cpy-ftf-NOS) was cloned into the XbaI and HindIII site of pMOG23 (Symons et al. supra), a binary plant vector pBIN19 derivative (Bevan M., supra) resulting in
10 plasmid pTP.

Transgenic plants, named TP plants, which contain the construct described were generated as described in Example 1.

15 3. Analysis of TP plants.

Screening of the transformants, using the TLC method as in Example 1 showed accumulation of fructan with the normal variation due to position effect. The fructans accumulating in these plants were further
20 studied by isolation of larger quantities (Livingstone III D.A., supra). This fructan was analysed by sizing on a FPLC Superose 6HR 10/30 column (Pharmacia) and fructans were detected in the void volume indicating a degree of polymerisation of >25.000 fructose units. Fructan produced by Streptococcus mutans fructosyltransferase similarly elutes in the
25 void volume of the Superose column. Similarly partial fructan degradation by acid degradation shows a characteristic pattern of hydrolysis products. The purified plant and bacterial fructans show identical degradation patterns on
30 TLC. Full acid hydrolysis followed by HPLC analysis on an Aminex HPX87C column at 85°C using water as eluent shows this fructan to be composed of fructose.

As in Example 1, senescing plants still contain fructans. Similarly seeds are fertile and the fructan producing property is inherited by following generation.
35

Example 3.

Expression of sacB in the apoplast

1. Selection of the genes.

To create plants which produce fructans, genes encoding proteins capable of producing fructans were selected. One of these genes, levansucrase encoded by the sacB gene (Steinmetz M. et al., supra) and described in Example 1, was used.

Since sucrose in tobacco plants is transported through the apoplast to the phloem, also the intercellular space (apoplast) is a preferred site for fructan accumulation. A gene encoding a protein capable of targeting itself to the apoplast was selected: the pathogenesis-related protein S, encoded by the pr-s gene (Cornelissen B.J.C. et al. Nature 321, 531-532 (1986). From this gene the export signal sequence was used.

2. Construction of 35S/pr-s-sacB/NOS in binary vector (pKT).

Plasmid pMOG29 has been described by Pen J., et al. (Bio/Technology 10, 292-296 (1992)). The pathogenesis-related protein S (pr-s) gene (Cornelissen B.J.C. et al., supra; Van Kan J.A.L. et al. Plant Mol. Biol. 12, 153-155 (1989)) has a signal sequence which can target proteins to the intercellular space, the apoplast (Pen J., et al., supra). The plasmid pMOG29 contains the 35S-promotor/pr-s-sequence/NOS-terminator construct. A pBluescript II SK (Stratagene) from which the internal BamHI site had been removed, by cutting with BamHI and filling in of the sticky ends with Klenow and religation was used for further cloning.

The 35S-promotor/pr-s-sequence/NOS-terminator construct was obtained by digestion with EcoRI and HindIII of pMOG29 and was cloned in this pBluescript into the corresponding EcoRI and HindIII site resulting in plasmid pPB1. Plasmid pPB1 was digested with BamHI and overhanging ends were filled in with Klenow. From plasmid pKK3 (Example 1.) the sacB gene was isolated by digestion with NcoI and PstI. These sites were blunted with Klenow. A blunt end ligation was performed resulting in plasmid pKR. The plasmid pKR encodes the in frame construct of the pr-s levansucrase fusion. The fusion

is in the correct reading frame which was confirmed by sequence analyses. The construct codes for the following amino acids sequence: MNFLKSFPFYAFLCFGQYFVAVTHARAS followed by the levansucrase protein sequence starting with methionine, alanine and the rest of the mature levansucrase protein (Steinmetz M. et al., supra). This construct was digested with XbaI and then partially digested with XhoI. The fragment containing the complete construct (35S/pr-s-sacB/NOS) was cloned into the XbaI and XhoI sites of pMOG23 (Symons et al. supra), a binary plant vector pBIN19 derivative (Bevan M., supra), resulting in plasmid pKT. Transgenic plants, named KT plants, which contain the construct described were generated as described in Example 1.

15 3. Analysis of KT plants.

Screening of the transformants, using the TLC method as in Example 1 showed accumulation of fructan with the normal variation due to position effect. The fructans accumulating in these plants were further studied by isolation of larger quantities (Livingstone III D.A., supra). Superose chromatography data and full acid hydrolysis followed by HPLC analysis as in Example 1 shows this fructan to be of high molecular weight and to be composed of fructose.

25 As in Example 1, senescing plants still contain fructans. Similarly seeds are fertile and the fructan producing property is inherited by following generation.

Example 4.

30 Expression of ftf gene in apoplast

1. Selection of the genes.

To create plants which produce fructans, genes encoding proteins capable of producing fructans were selected. Another gene, fructosyltransferase encoded by the ftf gene (Shiroza T. and Kuramitsu H.K. et al., supra) and introduced in Example 2, was used. Since sucrose in tobacco plants is transported through the apoplast to the phloem, also the intercellular space (apoplast) is a preferred site

for fructan accumulation. A gene encoding a protein capable of targeting itself to the apoplast was selected: the pathogenesis-related protein S, encoded by the pr-s gene (Cornelissen B.J.C. et al. supra). From this gene the signal sequence was used.

2. Construction of 35S/pr-s-ftf/NOS in binary vector (pTT)

Plasmid pPB1 (Example 3.) was digested with BamHI and overhanging ends were filled in with Klenow. From plasmid pTK1 (Example 2.) the ftf gene was isolated by digestion with NcoI and PstI. These sites were blunted with Klenow. A blunt end ligation was performed resulting in plasmid pTR. The plasmid pTR encodes the in frame construct of the pr-s fructosyltransferase fusion. The fusion is in the correct reading frame which was confirmed by sequence analyses. The construct encodes for the following amino acids sequence: MNFLKSFPFYAFLCFGQYFVAVTHARAS and fructosyltransferase protein sequence starting with the methionine of the mature fructosyltransferase (Shiroza T., supra)

This construct was digested with XbaI and HindIII. The fragment containing the complete construct (35S/pr-s-ftf/NOS) was cloned into the XbaI and HindIII site of pMOG23 (Symons et al., supra), a binary plant vector pBIN19 derivative (Bevan M., supra), resulting in plasmid pTT. Transgenic plants, named TT plants, which contain the construct described were generated as described in Example 1.

3. Analysis of TT plants.

Screening of the transformants, using the TLC method as in Example 1 showed accumulation of fructan with the normal variation due to position effect.

The fructans accumulating in these plants were further studied by isolation of larger quantities (Livingstone III D.A., supra). Superose chromatography data and full acid-hydrolysis followed by HPLC analysis as in Example 1 shows this fructan to be of high molecular weight and to be composed of fructose.

As in Example 1, senescing plants still contain fructans. Similarly seeds are fertile and the fructan producing property is inherited by following generation.

5 **Example 5.**

Expression of sacB in the cytoplasm

1. Selection of the genes.

To create plants which produce fructans, genes encoding proteins capable of producing fructans were selected. One of these genes, levansucrase encoded by the sacB gene (Steinmetz M. et al., supra) and described in Example 1, was used.

Since sucrose in plant cells is synthesized in the cytoplasm, the cytoplasm is a preferred site for fructan accumulation. Since nuclear-encoded proteins are made in the cytoplasm no targeting sequence is needed.

2. Construction of 35S-sacB-NOS in binary vector (pKZ)

Plasmid pPA2 (Example 1) was digested with NcoI and BamHI and the vector-containing fragment was isolated. From plasmid pKK3 (Example 1) the sacB gene was isolated as a with NcoI and BamHI fragment. Both fragments were ligated resulting in plasmid pKX. The resulting plasmid pKX encodes the construct of mature levansucrase. The construct is correct which was confirmed by sequence analyses.

This construct was digested with XbaI and XhoI. The fragment containing the complete construct (35S-sacB-NOS) was cloned into the XbaI and XhoI site of pMOG23 (-Symons et al., supra), a binary plant vector pBIN19 derivative (Bevan M., supra) resulting in plasmid pKZ.

Transgenic plants, named KT plants, which contain the construct described were generated as described in Example 1.

35 3. Analysis of KT plants.

Screening of the transformants, using the TLC method as in Example 1 showed accumulation of fructan with the normal variation due to position effect.

The fructans accumulating in these plants were further studied by isolation of larger quantities (Livingstone III D.A., supra). Superose chromatography data and full acid hydrolysis followed by HPLC analysis as in Example 1 shows this fructan to be of high molecular weight and to be composed of fructose.

As in Example 1, senescing plants still contain fructans. Similarly seeds are fertile and the fructan producing property is inherited by following generation.

Example 6.

Expression of ftf gene in cytoplasm

1. Selection of the genes.

To create plants which produce fructans, genes encoding proteins capable of producing fructans were selected. Another gene, fructosyltransferase encoded by the ftf gene (Shiroza T. and Kuramitsu H.K., supra) and introduced in Example 2, was used. Since sucrose in plant cells is synthesized in the cytoplasm, the cytoplasm is a preferred site for fructan accumulation. Since nuclear-encoded proteins are made in the cytoplasm no targeting sequence is needed.

2. Construction of 35S-ftf-NOS in binary vector (pTZ)

Plasmid pPA2 (Example 1) was digested with NcoI and BamHI and the vector-containing fragment was isolated. From plasmid pTK1 (Example 2) the ftf gene was isolated as a NcoI and BamHI fragment. Then both fragments were ligated resulting in plasmid pTX. The resulting plasmid pTX encodes the construct of mature fructosyltransferase. The construct is correct which was confirmed by sequence analyses.

This construct was digested with XbaI and HindIII. The fragment containing the complete construct (35S-ftf-NOS) was cloned into the XbaI and HindIII site of pMOG23 (Symons et al., supra), a binary plant vector pBIN19 derivative (Bevan M., supra), resulting in plasmid pTZ.

Transgenic plants, named TZ plants, which contain the construct described were generated as described in Example 1.

5 3. Analysis of TZ plants.

Screening of the transformants, using the TLC method as in Example 1 showed accumulation of fructan with the normal variation due to position effect.

10 The fructans accumulating in these plants were further studied by isolation of larger quantities (Livingstone III D.A., supra). Superose chromatography data and full acid hydrolysis followed by HPLC analysis as in Example 1 shows this fructan to be of high molecular weight and to be composed of fructose.

15 As in Example 1, senescing plants still contain fructans. Similarly seeds are fertile and the fructan producing property is inherited by following generation.

EXAMPLE 7.

20 Acquired properties of transgenic tobacco plants harbouring fructosyltransferase constructs.

1. General properties

The transgenic tobacco plant lines harbouring the different fructosyltransferase constructs as described in
25 examples 1-6 all accumulate fructan molecules. The level of fructan accumulation differed between individual transformants probably due to the effect of the integration site of the transgene on gene expression (position effect) and between the six different constructs used. Using Southern
30 hybridization analysis (Sambrook, et al supra) the number of DNA copies integrated into the individual plant genomes was determined. The number of copies integrations varied between 1-8, but most plant contained only one or two copies of the construct. The degree of polymerisation of the fructans is
35 up to 25.000 or more fructose units. Fructans are accumulated in all organs tested, including leaves, stem, roots and in the seeds. The property was transmitted to the offspring in an Mendelian manner.

The identity of the fructans was confirmed by sizing, hydrolysis and by proton nuclear magnetic resonance spectroscopy as described in the above examples. In addition antibodies directed against fructofuranosyl linkages (Hall, B. et al. 1990, Mol.Immunology 27: 351-361; Hall, B. et al. 1992, The Journal of Immunology 149: 1605-1612) reacted with the fructans produced in the transgenic plants. For this investigation fructans were spotted on nitrocellulose filter paper and immobilized by a 15 minutes treatment at 120°C. The filters were incubated with the specific mouse antibody (Hall et al., supra) and the binding of antibodies to the fructan detected with an alkaline phosphatase conjugated goat second antibody directed against the non-specific parts of the mouse antibody.

Fructans are present throughout the life cycle of the plant. Once synthesized the fructan is stable. If there is fructan turnover at all this level will be very low. In senescent dehydrated leaves fructans are still present in amounts comparable to levels in mature leaves. The fructan content in the transgenic tobacco plants is in the range of 3-8% of the dry weight in mature leaves. The levels of the other soluble carbohydrates (glucose, sucrose and fructose) were quantitated using HPLC chromatography as described and were found to be comparable to the untransformed wildtype plants.

2. Phenotypic characterisation

Under normal growth conditions no difference in growth or morphology was observed between the untransformed control tobacco plants and the fructan accumulating transgenic plants which harbour the *S. mutans* *ftf* gene in the three different configurations mentioned in examples 2, 4 and 6 (TP, TZ and TT lines). This is also true for plants harbouring the *B. subtilis* *sacB* gene fused to the *cpy* targeting signal (KP plants, example 1). Plant expressing the *B. subtilis* *sacB* gene in the cytoplasm or apoplast (KZ, KT, examples 3 and 5) show necrotic lesions in mature leaves and stems. The severity of the phenotype correlates with the

level of fructans accumulated. These plants do produce fertile seeds.

EXAMPLE 8

5 Enhanced performance under stress conditions of transgenic tobacco plants harbouring fructosyltransferase constructs.

The performance under stress conditions of representative fructan accumulating tobacco lines harbouring the cpy-sacB construct (KP, example 1) was compared with un-
10 transformed control plants.

1. Drought stress

Drought stress was induced in series of plants by the controlled addition to plants grown on vermiculite of
15 Poly Ethylene Glycol with a mean Molecular Weight of 10.000 dalton (PEG10.000). A range of drought stress levels was induced up to 20% PEG10.000 by application of PEG solutions. Fresh weight and dry weight was determined as well as growth rate (in cm/day). The drought stressed fructan accumulating
20 KP plants grew faster and gave significantly higher yields than similarly stressed non-transformed plants. When compared to similarly stressed non-transformed plants the transgenic KP plant fresh weight increased in a typical case up to 19% and dry weight up to 32%.

25

2. Light stress conditions

Yield characteristics under suboptimal light conditions were compared between representative transgenic KP plants and non-transformed plant. The KP plants grown
30 under low light (5000 lux) were directly compared to control (non-transformed) plants. The fructan accumulating KP plants grew faster and gave significantly higher yields than non-transformed plants. Both fresh weight and the dry weight of the KP plants was 18% higher than in non-transformed plants.
35 The KP produced more leaves than the non-transformed plants.

High light intensities, especially when combined with suboptimal growth conditions also results in severe stress in most plant species. Fructan accumulating plants

may under such conditions perform better than non-transformed plants. For example fructans function as a additional carbohydrate sink and in this way may protect against light induced damage to plants.

5

3. Cold stress

Growth characteristics of transgenic KP plants and non-transformed plants were compared under cold stress conditions. When grown under controlled condition at 12°C the KP plants gave over 19% higher yields in terms of fresh weight and dry weight than non-transformed plants.

The stress conditions named under 1), 2) and 3) induced a more than four-fold increase in the level of fructan in the the transgenic KP plants compared to unstressed KP plants. The accumulation of fructans results in improved growth performance, the degree of fructan accumulation being correlated with the degree of growth enhancement.

Better performance under stress conditions may not be limited to the specific examples given under 1), 2) and 3) but may occur under a wide range of unfavourable environmental conditions. Improved performance is not limited to increased growth rates or increases in fresh and dry weight but may include other economically important physiological aspect both with respect to pre- and post-harvest properties.

In addition to improved performance under abiotic stresses as the ones mentioned above, the fructan accumulating plants may also provide the plants with enhanced resistance against biotic stresses, like diseases and pests (Farrar, in: Pests and pathogens: plant response to foliar attack, p. 107-127. Ed. P. G. Ayres, BIOS Publishers 1992). Fructan accumulation may result in metabolic or structural modifications in plants resulting in reduced sensitivity to diseases and pests. Examples of such modifications include but are not limited to changes in digestibility, texture, flavour and altered levels of other plant primary and secondary metabolic products.

Plants with natural or induced modified carbohydrate partitioning patterns may be preferred targets for improved performance under stress conditions due, for example, to further increases in fructan accumulation. Such plants include but are not limited to natural mutants in starch and sucrose metabolism and plants in which starch and sucrose metabolism have been modified by molecular and genetic techniques for example as described in Sonnewald and Will-mitzer, Plant Physiology 99, 1267-1270, 1992.

EXAMPLE 9

General applicability of the technology

1. The construct

To demonstrate the general applicability of the technology use was made the cpy-sacB construct as described in example 1. Other genes of any origin that encode polypeptides able to synthesize fructan, minimally consisting of two linked fructose units, can be used as well. In addition, the effectiveness of the introduced constructs in terms of its ability to direct fructan accumulation in transgenic plants with respect to level, timing and location can be modified by using regulatory signals which include but are not limited to constitutive promoters, organ specific promoters, developmentally regulated promoters, polyadenylation signals, translational enhancers, transcriptional enhancers etc. In addition the cellular localization of the polypeptide can be directed by using different cellular targeting signals which include but are not limited to vacuolar and apoplastic targeting signals of any origin. After introduction into plants of any combination of the above mentioned elements significant quantities of fructans can accumulate in vegetative organs which include leaves, roots and stems as well as derived organs which include but are not limited to tubers, storage roots, fruits as well as seeds.

2. Application in different crop species

To demonstrate the general applicability of the technology the construct cpy-sacB as described in example 1

was introduced into crop species like potato (Solanum tuberosum L.) of which the transformation can be but does not need to be carried out as described in Visser, Plant Tissue Culture Manual B5, 1-9, Kluwer Academic Publishers, 1991.

- 5 The resulting transgenic plants accumulated fructans in every organ throughout development. Levels in leaves and tuber can be over 9% of the fresh weight. In addition the same construct was introduced into beet (Beta vulgaris L.) which can be but does not need to be transformed as described by D'Halluin et al. Biotechnology 10, 309-314 (1992).
10 The resulting transgenic beet plants accumulated significant levels of fructan with a degree of polymerization up to 25.000 or more in for example their leaves and storage roots. The same cpy-sacB construct was introduced into
15 fodder brassica (Brassica napus L.) which can be but does not need to be transformed as described by de Block et al. Plant Physiol 91, 694-701 (1989). The resulting transgenic fodder brassica plants accumulated significant levels of fructans with a degree of polymerization up to 25.000 or
20 more in for example their leaves and storage organs.

In addition plant species that can be modified to accumulate fructans include but are not limited to maize (Zea mays L.), wheat (Triticum aestivum L.), barley (Hordeum vulgare L.), rice (Oryza sativa L.) soybean (Glycin max L.),
25 pea (Pisum sativum L.), bean (Phaseolus vulgaris L.) chicory (Cichorium intybus L.), sugarcane (Saccharum officinarum L.), yam (Dioscorea esculenta L.), cassave (Manihot esculenta L.) and grasses (e.g. Lolium spp., Poa spp. and Festuca spp.).

- 30 Plants with natural or induced modified carbohydrate partitioning patterns may be preferred targets for modification of fructan metabolism, especially increased levels of fructan accumulation. Such plants include but are not limited to natural mutants in starch and sucrose metabolism and plants in which starch and sucrose metabolism have
35 been modified by molecular and genetic techniques for example as described in Sonnewald and Willmitzer, Plant Physiology 99, 1267-1270, 1992.

CLAIMS

1. Method for obtaining transgenic plants showing
5 a modified fructan pattern as compared to non-transformed
plants, comprising the steps of:
- a) preparing a DNA construct comprising one or
more fructosyltransferase genes, or modified versions there-
of, being operably linked to a promotor sequence active in
10 plants and a terminator sequence active in plants;
- b) transforming a plant cell with the construct;
and
- c) regenerating a transgenic plant from the trans-
formed plant cell.
- 15 2. Method as claimed in claim 1, characterized in
that, the DNA construct also comprises a targeting sequence
upstream from the fructosyltransferase gene.
3. Method as claimed in claim 1 or 2, characteri-
zed in that, the 5' untranslated region of the fructosyl-
20 transferase gene is modified such that the expression of the
fructosyltransferase is not negatively affected.
4. Method as claimed in claim 1, 2 or 3, character-
ized in that, the fructosyltransferase gene originates from
a micro-organism.
- 25 5. Method as claimed in claim 4, characterized in
that, the fructosyltransferase gene selected from the group
consisting of the sacB gene of Bacillus subtilis, the ftf
gene of Streptococcus mutans.
6. Method as claimed in claim 1 or 2, characteri-
30 zed in that, the fructosyltransferase gene originates from a
plant.
7. Method as claimed in any one of the claims 2-6,
characterized in that, the targeting sequence is selected
from the group consisting of the signal sequence and vacuo-
35 lar targeting sequence of the carboxypeptidase Y (cpv) gene,
the signal sequence and apoplasmatic targeting sequence of
the pathogenesis-related protein S gene.

8. Method as claimed in any one of the claims 1-7, characterized in that, the terminator sequence is a nopaline synthase gene terminator sequence.

9. Method as claimed in any one of the claims 1-8,
5 characterized in that, the promotor sequence is a constitutive or regulable promotor sequence.

10. Method as claimed in any one of the claims 1-8, characterized in that, the promotor sequence is selected from the group consisting of the 35S Cauliflower Mosaic
10 Virus promotor sequence, the potato granule-bound starch-synthase promoter.

11. Method as claimed in any one of the claims 1-10, characterized in that, the DNA construct also comprises a translational enhancer sequence upstream from the fructosyltransferase gene and downstream from the promoter sequence.
15

12. Method as claimed in claim 11, characterized in that, the translational enhancer sequence is a Alfalfa Mosaic Virus RNA4 translational enhancer.

13. DNA construct for the production of transgenic plants, comprising one or more fructosyltransferase genes, or modified versions thereof, being operably linked to a plant specific promotor sequence and a plant specific terminator sequence.
20

14. DNA construct as claimed in claim 13, characterized in that, the 5' untranslated region of which is modified such that the expression of the fructosyltransferase is not negatively affected.
25

15. DNA construct as claimed in claim 13 or 14, also comprising a targeting sequence upstream from the fructosyltransferase gene.
30

16. Transgenic plant cell comprising a DNA construct as claimed in claim 13, 14 or 15.

17. Transgenic plant produced according to the
35 method of any one of the claims 1-12.

18. Transgenic plant produced according to the method as claimed in claims 1-12 tolerant to abiotic and/or biotic stresses.

19. Transgenic plant tissue, such as fruit, stem, root, tuber, seed of a plant as claimed in claim 17 or 18.

20. Use of a transgenic plant as claimed in claim 18 for culturing under conditions of abiotic and/or biotic stress.

21. Use of the transgenic plant as claimed in claim 17 or the transgenic plant tissue as claimed in claim 19 for the production of fructans.

22. Use of the transgenic plant as claimed in claim 17 or the transgenic plant tissue as claimed in claim 19 as a food stuff.

23. Use of the transgenic plant as claimed in claim 17 or the transgenic plant tissue as claimed in claim 19 as feed for animals.

24. Fructans isolated from a transgenic plant as claimed in claim 17 or a plant tissue as claimed in claim 19.

25. Use of the fructans as claimed in claim 24 in food and non-food products and in industrial processes

26. Use of the fructans as claimed in claim 25 in low-calory food products.

27. Transgenic seed obtainable from a transgenic plant as claimed in claim 17 or 18.

28. Reproducible structure obtainable from a transgenic plant as claimed in claim 17 or 18.





FIG. 1



A=AccI
B=BamHI
E=EcoRI
H=HindIII
K=KpnI
N=NcoI
P=PstI
S=SphI
X=XhoI
Xho=XhoI




FIG. 2

FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7





FIG. 8

Degradation of Tobacco Levan KP12 **200792**



INTERNATIONAL SEARCH REPORT

Int. .onal Application No

PCT/NL 93/00279

A. CLASSIFICATION OF SUBJECT MATTER

IPC 5 C12N15/82 C12N15/54 C12N5/10 A01H5/00 A23L1/00
A23L1/09 A23L1/307 A23K1/00 C08B37/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 5 C12N A01H A23L A23K C08B

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	BIOCHEMICAL SOCIETY TRANSACTIONS vol. 19, no. 3, 1991 pages 569 - 572 JOHN, P. 'Fructan quality and fructan synthesis' see the whole document	24-26
A	WO,A,89 12386 (CALGENE) 28 December 1989 cited in the application see example 4	1-28
E	WO,A,94 04692 (INSTITUT FÜR GENBIOLOGISCHE FORSCHUNG BERLIN) 3 March 1994 see the whole document	1,2,4,9, 10,13, 15-21, 24,28

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- 'A' document defining the general state of the art which is not considered to be of particular relevance
'E' earlier document but published on or after the international filing date
'L' document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
'O' document referring to an oral disclosure, use, exhibition or other means
'P' document published prior to the international filing date but later than the priority date claimed

'T' later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

'X' document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

'Y' document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

'&' document member of the same patent family

Date of the actual completion of the international search

13 April 1994

Date of mailing of the international search report

27-04-94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Maddox, A

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/NL 93/00279

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-8912386	28-12-89	AU-A- 3852089	12-01-90
WO-A-9404692	03-03-94	DE-A- 4227061	17-02-94